## In the Claims:

Please cancel Claims 2, 3, 5, 6, 8, and 9 and amend Claims 1, 4 and 7 as follows:

1. (Currently Amended) A DNA microarray comprising:

a set of features on a substrate, each of the features including <u>multiple copies of single</u> stranded DNA probes <u>of common sequence</u>, <u>and</u>

features including positive control probes being included in the set of features, the probes features for the positive control probes controls being arranged in a pattern on the microarray such that the features having the positive control probes create a symbol recognizable to a human being through visual observation when illuminated, so that whether an event of interest has occurred can be determined by hybridizing fluorescently tagged nucleic acids from a sample to the microarray, illuminating the microarray, and observing the presence or absence of the symbol visual pattern.

## 2.-3. (Cancelled)

4. (Currently Amended) A method for building designing a polynucleotide microarray comprising the steps of:

selecting a set of features, each feature including <u>a plurality of</u> polynucleotide probes <u>of identical nucleotide sequence</u> for detecting an event of interest, some of the features <u>including probes designed to serve as being</u> positive controls; and

positive controls when illuminated form a pattern symbol recognizable to a human being through visual observation if the positive control features including positive control probes fluoresce,

wherein the set of features provides a polynucleotide microarray.

## 5.-6. (Cancelled)

7. (Currently Amended) A method for detecting whether an event of interest in a biological experiment has occurred comprising the steps of:

providing a DNA microarray comprising a set of features, each <u>feature</u> including <u>a</u> <u>plurality of</u> single stranded DNA probes <u>of the same sequence</u>, for detecting the event of <u>interest</u>, the microarray including features intended to serve as positive controls <u>indicating</u> that the event of interest has occurred, the features for positive controls being arranged in a pattern <u>forming a character</u> recognizable to a human being through visual observation;

hybridizing nucleic acids from a sample to the microarray; and observing the presence or absence of the visual pattern to determine if the event of interest has occurred.

8.-9. (Cancelled)